

Exercises to the lecture  
Complexity Theory  
Sheet 9

Prof. Dr. Roland Meyer

M.Sc. Peter Chini

Delivery until 15.01.2018 at 18h

**Exercise 9.1** (Modeling via Polynomials)

We use the polynomial approach to describe the solutions to the following problem:

**k-PATH****Input:** A directed graph  $G = (V, E)$  and  $k \in \mathbb{N}$ .**Parameter:**  $k$ .**Question:** Is there a path of  $k$  vertices in  $G$ ?

To come up with the right polynomial, we need some preliminaries: A *walk* is a sequence of  $k$  vertices  $v_1, \dots, v_k \in V$ , with potential repetition. A *labeling* is a bijective function  $\ell : [1..k] \rightarrow [1..k]$ . A *labeled walk* is a pair  $(W, \ell)$ , where  $W$  is a walk and  $\ell$  is a labeling. Intuitively, every (visit of a) vertex  $v$  in  $W$  gets a unique label from  $\ell$ .

Our variables are  $X = \{x_{u,v} \mid (u,v) \in E\}$  and  $Y = \{y_{v,j} \mid v \in V, j \in [1..k]\}$ . Let  $(W, \ell)$  be a labeled walk with  $W = v_1, \dots, v_k$ , then we define the monomial:

$$\text{mon}_{W,\ell}(X, Y) = \prod_{i=1}^{k-1} x_{v_i, v_{i+1}} \cdot \prod_{i=1}^k y_{v_i, \ell(i)}.$$

Note that the monomial models precisely the labeled walk in terms of variables.

The instance polynomial is given by the sum of all such monomials:

$$P(X, Y) = \sum_{\substack{\text{walk} \\ W=v_1, \dots, v_k}} \sum_{\ell \text{ a labeling}} \text{mon}_{W,\ell}(X, Y).$$

We show that  $P(X, Y)$  does not vanish modulo 2 if and only if  $G$  contains a  $k$ -path.

- a) We define a function  $T$  between labeled walks. Let  $(W, \ell)$  be a labeled walk with  $W = v_1, \dots, v_k$ . If  $W$  is not a path then there is a smallest pair of numbers  $(a, b)$  such that  $a < b$  and  $v_a = v_b$ . Define the labeling  $\ell'$  as follows:

$$\ell'(x) = \begin{cases} \ell(b), & x = a \\ \ell(a), & x = b \\ \ell(x), & \text{otherwise.} \end{cases}$$

The function  $T$  maps a labeled walk  $(W, \ell)$  to the labeled walk  $(W, \ell')$  if  $W$  is not a path. If  $W$  is a path,  $T$  is the identity map.

Prove that  $\text{mon}_{W,\ell} = \text{mon}_{T(W,\ell)}$  for each labeled walk  $(W, \ell)$ .

b) Show that  $P(X, Y)$  takes the following form modulo 2:

$$P(X, Y) = \sum_{\substack{\text{path} \\ W=v_1, \dots, v_k}} \sum_{\ell \text{ a labeling}} \text{mon}_{W, \ell}(X, Y).$$

c) Conclude that  $P(X, Y)$  does not vanish modulo 2 if and only if  $G$  contains a  $k$ -path.

**Exercise 9.2** (Zeta and Möbius Transform)

Prove the following equivalences:

a)  $\zeta = \sigma\mu\sigma$ , and

b)  $\mu = \sigma\zeta\sigma$ .

*Hint:* Recall that  $\sigma\sigma = id$ .

**Exercise 9.3** (Cover Product)

Let  $V$  be a finite set and  $f, g : \mathcal{P}(V) \rightarrow \mathbb{Z}$  functions. We define the *cover product* of  $f$  and  $g$  to be the function  $f *_c g : \mathcal{P}(V) \rightarrow \mathbb{Z}$  with

$$(f *_c g)(X) = \sum_{\substack{A, B \subseteq X \\ A \cup B = X}} f(A) \cdot g(B).$$

Show that  $\zeta(f *_c g) = (\zeta f) \cdot (\zeta g)$

**Delivery until 15.01.2018 at 18h into the box next to 343**